

An algebraic characterization of singular quasi-bi-Hamiltonian systems

Rolando Alvarado F^a., Maximo Agüero G^b.

^a*Centro de Estudios Multidisciplinarios,
Universidad Autonoma de Zacatecas
Apartado Postal 597 - C, Zacatecas, Zac. Mexico.*

Email: ralva@cantera.reduaz.mx

^b*Universidad Autonoma del Estado de Mexico,
Facultad de Ciencias, Instituto Literario 100,
Toluca 50000, Edo. Mex., Mexico.*

Email: mag@coatepec.uaemex.mx

February 6, 2008

Abstract

In this paper we prove an algebraic criterion which characterizes singular quasi-bi-hamiltonian structures constructed on the lines of a general, simple, new formal procedure proposed by the authors. This procedure shows that for the definition of a quasi-bi-hamiltonian system the requirement of non-singular Poisson tensors, contained in the original definition by Brouzet et al., is not essential. Besides, it is incidentally shown that one method of constructing Poisson tensors available in the literature is a particular case of ours. We present 2 examples.

1 Introduction.

The use of Hamiltonian methods in physics is as old as mathematical-physics and can be traced back to the works of Euler and Lagrange in point and con-

tinuum mechanics. Recently these methods, which we may divide in symplectic and Poisson, have been developed in the works of Kostant[11], Kirillov[9], Lichnerowicz[8], Guillemin[13], Soriau[12], Weinstein[10] and others in various fields which cover from geometric quantization to control theory[15].

One current interest is now to construct Hamiltonian theories without the help of a Lagrangian function[1], mainly because sometimes this way of constructing the theory is not available (just remember the celebrated Dirac theory of constraints). We mean, the hessian matrix of the Lagrangian function has a rank less than the dimension of the configuration space, hence, a Hamiltonian formulation does not seem available. Of course, the equations of motion in the usual coordinates of the Lagrangian are always, or must be always, available. So, they can be considered as the starting point in the construction of a Hamiltonian formulation. Clearly, if we have at hand a given Hamiltonian formulation, it is interesting to know if we can construct another one over this. However we must be clear as to what kind of structure we wish to get on a manifold. It is possible to construct a symplectic structure, which in the local coordinates of the symplectic manifold give rise to the Lagrange brackets, or we may try to construct a Poisson structure and, of course, the Poisson brackets in the local coordinates of the Poisson manifold; which in the degenerate case cannot be used to deduce, by inversion, the symplectic structure. In this paper we will restrict ourselves to the discussion of a new and simply method to construct singular quasi-bi-Hamiltonian structures; in the sense in which Brouzet et al. defined this concept in their interesting paper[2]; with the help of an algebraic criterion deduced utilizing a decomposable Poisson tensor. The main difference with the procedure of Brouzet et al. is that we do not require the existence of any symplectic structure on the manifold, we require only Poisson tensors which we allow to be singular (degenerate); besides, our treatment is useful for any dimension of the underlying manifold. Hence, the definition of a quasi-bi-Hamiltonian structure is independent of any underlying symplectic structure. For this reason we call the quasi-bi-Hamiltonian systems constructed "singular". We will achieve this goal starting directly from a known Poisson tensor, in order to erect another one on this basis. As we shall see, the method provides a technique for defining the Hamiltonians for both structures, so in our approach the only data that we need is a Poisson tensor. This is in accordance with the usual procedures followed by some professional constructors of Poisson structures[1][5], with just one difference: some of them use a first order differential condition on the Hamiltonian function (it is a constant of the

motion) but we use a second order condition, the Hamiltonian is the solution of a second order differential equation, and we may consider that the usual first order condition is a particular case of our condition.

In the next section (2) we give some useful definitions for the full comprehension of the paper. In section (3) we give the main results, contained in theorems (1) and (2).

In section (4) we give a brief introductory discussion of the Jacobi structure from the point of view of our general methodology, in section (5) we give 3 examples of the method and in the last section (6) we give the conclusions, which try to illuminate the full discussion in the article.

2 The definition of Poisson structures.

Let M be a smooth manifold and $C^\infty(M, \mathbb{R})$ the ring of all real valued, infinitely differentiable, functions on M [15]. A Poisson structure on M is given by a bilinear operation: $\{, \}$ on $C^\infty(M, \mathbb{R})$ such that the maps: $X_H = \{H, *\}$, $L_H = \{*, H\}$ are derivations, this operation is known as the "Poisson bracket". A manifold endowed with a Poisson bracket on C^∞ is called a Poisson manifold and we will denote it by the pair: $O = \langle M, \{, \} \rangle$.

We may understand by a Poisson structure on a manifold the explicit definition of a Poisson bracket, or the definition of a contravariant antisymmetric 2-tensor defined at all the points of M . In local cosymplectic coordinates this tensor is:

$$\{F, G\} = J^{ij}(x) \frac{\partial F}{\partial x^i} \frac{\partial G}{\partial x^j}. \quad (1)$$

Usually, to specify the 2-tensor we use the fundamental Poisson brackets of the local coordinates: $J^{ij}(x) = \{x^i, x^j\}$. The Jacobi identity gives us a first order partial differential equation which the 2-tensor must satisfy, but we will not display it here.

The tensor defines an isomorphism between the cotangent bundle T^*M and the tangent bundle TM if, and only if, it is non-degenerate. However, we will not consider this as an essential condition in the definition of the Poisson structure, because, as we will see, it is not necessary for the construction of the quasi-bi-Hamiltonian structure. If we use the 2-tensor, we will use the pair: $O = \langle M, J \rangle$ to express in a coordinate-free manner the Poisson manifold.

DEFINITION 1.- We will say that we have given a Hamiltonian structure on the Poisson manifold O when we give the scalar generator of the dynamics $H \in C^\infty(M, \mathbb{R})$, which is known as the Hamiltonian of the Hamiltonian structure. We will denote this by the triplet: $K_H = \langle O, H \rangle = \langle M, \{, \}, H \rangle$.

To define the dynamics in the local coordinates of O we need only the Poisson bracket and the Hamiltonian, because with these elements we may define a dynamics, locally or globally, as the integral curves of the Hamiltonian vector field given by:

$$X_H = \{x^i, x^j\} \frac{\partial H}{\partial x^i} \frac{\partial}{\partial x^j}$$

in local coordinates. So, in all the constructions of a Poisson structure it is important to define the set of all possible Hamiltonians.

DEFINITION 2.-[4] We will say that we have a bi-Hamiltonian structure if, and only if, given a Hamiltonian structure: $K_1 = \langle M, \{, \}_1, H_1 \rangle$, it is possible to construct another different Hamiltonian structure: $K_2 = \langle M, \{, \}_2, H_2 \rangle$ such that: $\{x^i, H_1\}_1 = \{x^i, H_2\}_2$. So, the dynamics accept two different formulations:

$$\frac{dx^i}{dt} = \{x^i, H_1\}_1 = \{x^i, H_2\}_2. \quad (2)$$

The attempt of construction of this kind of formulations, even in the singular (degenerate) case, have led to the development of a series of techniques for its effective realization[4].

The previous definitions are standard in the theory of bi-Hamiltonian systems; now we must pass to the quasi-bi-Hamiltonian case of Brouzet et al.[2] but, as we wish to treat the singular case, we give a modified definition of a quasi-bi-Hamiltonian system in such a way that the singularities are not important:

DEFINITION 3.- Let K_H be a Hamiltonian system. We say that it admits a quasi-bi-Hamiltonian structure if, and only if:

(i).- There exists on M a Poisson tensor $J_\wedge$ compatible with J , i. e. its Schouten bracket (for the definition and properties of the Schouten bracket we use the book by Marsden and Ratiu[15]) commutes.

(ii).- There exists a non-vanishing function $\rho \in C^\infty(M, \mathbb{R})$ such that: $\rho (dH \rfloor J)$ is a globally Hamiltonian vector field.

Here $\rfloor$ denotes the contraction operation on a tensor field. We shall denote the quasi-bi-Hamiltonian system with:

$QBH = \langle K_H, J_\wedge, \rho \rangle = \langle M, J, H, J_\wedge, \rho \rangle$. We will see later how to get the second Hamiltonian.

If $\rho(dH \rfloor J)$ is globally Hamiltonian[6] for J , i. e. $(\exists F) : \rho(dH \rfloor J) = X_F$, then QBH is called "exact". As we shall see later, the function F is the solution of a partial differential equation of first order, and we shall understand the solutions to this equation in a local sense, as given by the usual theorems of existence[17]. In this way we can make the following difference: if the solution is available in closed form the vector field $\rho(dH \rfloor J)$ is globally Hamiltonian, but if the solutions are only available through the theorems of existence, usually power series which converge in some open disk, we shall call the vector field $\rho(dH \rfloor J)$ a locally Hamiltonian vector field¹.

The definitions in terms of symplectic structures, hence with the recursion operator at hand (Nijenhuis operator), were given by R. Brouzet et al.[2] and can be deduced from (i) and (ii) if both Poisson tensors are non-degenerate, because in such a case we have: $\omega = -J^{-1}$, $\omega_\wedge = -J_\wedge^{-1}$ hence the Nijenhuis operator is: $-J_\wedge \circ \omega$. Definition (3) can be changed by that of Brouzet et al. just replacing the Schouten bracket with the Nijenhuis torsion for the test of the compatibility of the symplectic structures (which is, of course, more complicated from the point of view of calculations than the relatively easier Schouten bracket). Hence, in this sense the extension of our definition covers that one of Brouzet et al., because even when the symplectic structure is not available, our definition has no problem. It is necessary to remark that the definition is introduced because it shows that the necessity of an inverse for the Poisson structure is not an important condition.

3 Compatibility of Poisson structures.

The notation in this section is as follows: $[\cdot, \cdot]$ denotes the Schouten bracket. We start with the well-known:

DEFINITION 4.-[3] A Poisson tensor, $\wp$, will be called decomposable (or

¹In the symplectic framework we call a Hamiltonian vector field a locally Hamiltonian vector field if, and only if, its contraction with the symplectic two-co-tensor is zero. In symbols, if Ω is the two-co-tensor and X the vector field we have: $d(X \rfloor \Omega) = 0$ and by Poincare lemma we have that locally: $X \rfloor \Omega = dH$ [see ref[15]. p.141]. Hence our introduced notion for the vector field $\rho(dH \rfloor J)$ is not arbitrary.

the Poisson tensor J_1 extendable if there is a Poisson tensor J_2 such that we may compose the tensor $\wp$) if, and only if, it is possible to write it as: $\wp = J_1 + J_2$ where each tensor J_1, J_2 is a Poisson tensor and: $[J_1, J_2] = 0$.

We will call the tensors which commute under the Schouten bracket: compatible tensors. As is well known[3], extensions of Poisson tensors can be used to study the problem of classification of solvable Lie algebras.

The methodology which we will use here to extend Poisson tensors is as follows: given a Poisson tensor of the form $J = X_1 \wedge X_2$, with the two vector fields: $X_1 = f_i \frac{\partial}{\partial x^i}$, $X_2 = g_i \frac{\partial}{\partial x^i}$ and a third vector field X_3 , we construct the new Poisson tensor J_γ as $X_\gamma \wedge X_3$, where $X_\gamma = \gamma \rfloor J$ and γ is a 1-co-tensor which we shall consider of the integrable (exact) form: $\gamma = dH$, with H a real valued function. Note that the choice of Poisson tensors is not general, however, this lack of generality is presented in many constructions of this kind.

So, if we follow the methodology just described to construct a new Poisson tensor over an old one, we only need to show that the new Poisson tensor commutes with the old one, and this procedure will give us the conditions to determine the vector field X_3 (the conditions obtained are independent of the representation used by the generators of the algebra.). Before proceed to the explicit constructions it is very important to see that in the methodology one crucial step is the choice of the form of two objects:

(A).-the Poisson tensor $J = X_1 \wedge X_2$ which is not the most general one, but however is the form used by Hojman (see ref.[1] p.669. In the next section we establish the explicit relation with the method by Hojman. Even in the case of J. Goedert[5] are required some restrictions, specifically the space dimension).

(B).- The choice of the vector field $X_H = dH \rfloor J$, as the contraction of the Poisson tensor.

These suppositions are independent in the sense that it is only in the lemma4 and the identity (7) that both are used to get a reduced set of conditions involving the vectors X_1, X_2, X_3 , in such a way that it is possible to know the third vector in terms of the first two, i. e. the second Poisson tensor with just the elements of the first one. We remark this because, as we shall see in the next section, with this choice we can re-construct the quasi-bi-Hamiltonian system in a very easy way.

We start with a few easy lemmas (in the development we shall suppose that the vector fields X_1, X_2, X_3 are linearly independent. There are no conditions on X_H unless otherwise stated):

LEMMA1.- The 2-contra-tensors $X_1 \wedge X_2$, $X_H \wedge X_3$ are Poisson tensors, respectively, if, and only if:

$$[X_1, X_2] = N_1 X_1 + N_2 X_2 \quad (3a)$$

$$[X_H, X_3] = A_1 X_H + A_2 X_3, \quad (3b)$$

where N_1, N_2, A_1, A_2 are arbitrary functions of the coordinates.

PROOF: the Schouten brackets for each tensor are:

$$X_1 \wedge [X_1, X_2] \wedge X_2 = 0$$

$$X_H \wedge [X_H, X_3] \wedge X_3 = 0$$

So, we can see that the conditions (3a-b) are sufficient conditions, because if they hold, then the Schouten bracket vanishes, as a simple calculation shows. To the opposite side: if the Schouten bracket is zero, then the conditions (3a-b) are the only solutions.●

LEMMA2.- X_H is an infinitesimal automorphism of the Poisson tensor $X_1 \wedge X_2$ if, and only if (to avoid any confusion: we use the notation X_H for convenience, it does not mean that the vector field is Hamiltonian):

$$[X_H, X_1] = -C_1 X_1 + B_2 X_2 \quad (4a)$$

$$[X_H, X_2] = C_1 X_1 + C_2 X_2 \quad (4b)$$

where C_1, C_2, B_2 are arbitrary functions of the coordinates.

PROOF:

The Lie derivative of the tensor $X_1 \wedge X_2$ with respect to the vector field X_H is, in terms of the Schouten bracket:

$$[X_H, X_1 \wedge X_2] = [X_H, X_1] \wedge X_2 + X_1 \wedge [X_H, X_2] = 0$$

clearly if X_H is an infinitesimal automorphism this Lie derivative must be zero. We can see that the conditions (4a-b) are sufficient, because if they hold then the Lie derivative is zero, as a substitution shows. They are necessary too because if the Schouten bracket vanishes, the only solution for it is through conditions (4a-b).●

LEMMA3.-Suppose that X_H is an infinitesimal automorphism of the tensor $X_1 \wedge X_2$, then this tensor and the tensor $X_H \wedge X_3$ are compatible tensors if, and only if:

$$[X_3, X_1] = D_1 X_H + D_2 X_2 \quad (5a)$$

$$[X_3, X_2] = E_1 X_H + E_2 X_2 \quad (5b)$$

PROOF: The Schouten bracket of the tensors is:

$$([X_H, X_1] \wedge X_2 + X_1 \wedge [X_H, X_2]) \wedge X_3 - X_H \wedge ([X_3, X_1] \wedge X_2 + X_1 \wedge [X_3, X_2]),$$

but by the lemma2 (which we suppose to hold, i. e. the relations 4a-b are valid) the first term is zero, hence we get the equation:

$$X_H \wedge ([X_3, X_1] \wedge X_2 + X_1 \wedge [X_3, X_2]) = 0.$$

So, the result (5a-b) is an immediate consequence, because if the conditions (5a-b) hold, the Schouten bracket vanishes, and if the Schouten bracket vanishes the only solution to the equation is through conditions (5a-b).•

THEOREM1.- Given four vectors X_1, X_2, X_3, X_H , then the tensors $X_1 \wedge X_2, X_H \wedge X_3$ are compatible Poisson tensors such that X_H is an infinitesimal automorphism of $X_1 \wedge X_2$ if, and only if, they form the basis of an algebra with the following commutation relations:

$$[X_1, X_2] = N_1 X_1 + N_2 X_2 \quad (6a)$$

$$[X_H, X_3] = A_1 X_H + A_2 X_3 \quad (6b)$$

$$[X_H, X_1] = -C_2 X_1 + B_2 X_2 \quad (6c)$$

$$[X_H, X_2] = C_1 X_1 + C_2 X_2 \quad (6d)$$

$$[X_3, X_1] = D_1 X_H + D_2 X_2 \quad (6e)$$

$$[X_3, X_2] = E_1 X_H + E_2 X_1 \quad (6f)$$

PROOF.- just use the lemmas (1-2-3).•

In this way the notions of compatibility of Poisson tensors and the property of one vector of being an infinitesimal automorphism of one Poisson tensor are algebraic concepts whose structure is contained in the relations (6a-6f).

This is a 4-dimensional algebra which we shall reduce to a 3-dimensional one with the help of the condition: $X_H = X_1(dH)X_2 - X_2(dH)X_1$. We must remark that this condition has not been used in the development at any point, so it is an independent condition. Next, we shall use the identity:

$$[X_H, X_i] = X_1(dH)[X_2, X_i] + X_2(dH)[X_1, X_i] - X_i X_1(dH)X_2 - X_i X_2(dH)X_1 \quad (7)$$

which follows from the derivation property of the commutator and the form which we have supposed for the vector field X_H . Note that the Lie bracket is not $C^\infty(M, \mathfrak{R})$ -bilinear and that the identity (7) change if we use another form for the infinitesimal automorphism.

LEMMA4.- If $X_H = X_1(dH)X_2 - X_2(dH)X_1$ the 4-dimensional algebra (6a-f) is reduced to the 3-dimensional algebra (of 3 linearly independent vectors):

$$\begin{aligned} [X_1, X_2] &= N_1 X_1 + N_2 X_2 \\ [X_3, X_1] &= (D_1 X_1(dH) + D_2) X_2 - D_1 X_2(dH) X_1 \\ [X_3, X_2] &= (E_2 - E_1 X_2(dH)) X_1 - E_1 X_1(dH) X_2 \end{aligned}$$

if we choose the functions:

$$\begin{aligned} C_1 &= X_2(dH)N_1 - X_2^2(dH) \\ C_2 &= X_1 X_2(dH) - X_1(dH)N_1 \\ B_1 &= -C_2 \\ B_2 &= \frac{X_1(dH)}{X_2(dH)}(X_1 X_2(dH) - X_1(dH)N_1) + \frac{X_2 X_1(dH)}{X_2(dH)} - X_1^2(dH) \\ N_2 &= \frac{X_1 X_2(dH)}{X_2(dH)} - \frac{X_1(dH)}{X_2(dH)}N_1 + \frac{X_2 X_1(dH)}{X_2(dH)} \\ A_1 &= \frac{X_1(dH)}{X_2(dH)}E_2 - X_2(dH)D_1 - X_1(dH)E_1 + \frac{X_3 X_2(dH)}{X_2(dH)} \\ A_2 &= -\left(X_2(dH)D_2 + \frac{(X_1(dH))^2}{X_2(dH)}E_2 + X_3 X_1(dH) + \frac{X_3 X_2(dH)X_1(dH)}{X_2(dH)}\right) \end{aligned}$$

leaving undetermined the remaining functions: D_1, D_2, E_1, E_2, N_1 .

PROOF.-The idea used here is quite simple: we shall try to satisfy identically the commutation relations (6b-c-d) with the help of the form which we use for the vector field X_H and an adequate selection of the arbitrary functions.

So, with this in mind, we use the identity (7) and the form of X_H to change the algebra (6a-f) to:

$$[X_1, X_2] = N_1 X_1 + N_2 X_2 \quad (8a)$$

$$X_1(dH)[X_2, X_3] - X_3 X_1(dH)X_2 + X_2(dH)[X_1, X_3] - X_3 X_2(dH)X_1 = A_2 X_2 + A_1(X_1(dH)X_2 - X_2(dH)X_1) \quad (8b)$$

$$X_1(dH)[X_2, X_1] - X_1^2(dH)X_2 - X_1 X_2(dH)X_1 = -C_2 X_1 + B_2 X_2 \quad (8c)$$

$$-X_2 X_1(dH)X_2 + X_2(dH)[X_1, X_2] - X_2^2(dH)X_1 = C_1 X_1 + C_2 X_2 \quad (8d)$$

$$[X_3, X_1] = D_1(X_1(dH)X_2 - X_2(dH)X_1) + D_2 X_2 \quad (8e)$$

$$[X_3, X_2] = E_1(X_1(dH)X_2 - X_2(dH)X_1) + E_2 X_1 \quad (8f)$$

Now we put (8a) in (8c) and (8d) and (8e), (8f) in (8b). From the substitution of (8a) in (8c) and (8d) we get the equations:

$$\begin{aligned} (X_1(dH)N_1 + C_2 - X_1 X_2(dH))X_1 + (X_1(dH)N_2 - X_1^2(dH) - B_2)X_2 &= 0 \\ (X_2(dH)N_1 - X_2^2(dH) - C_1)X_1 + (X_2(dH)N_2 - C_2 - X_2 X_1(dH))X_2 &= 0 \end{aligned}$$

so, by the linear independence of the vector fields at each point of the manifold, we get the equations:

$$X_1(dH)N_1 + C_2 - X_1 X_2(dH) = 0 \quad (9a)$$

$$X_1(dH)N_2 - B_2 - X_1^2(dH) = 0 \quad (9b)$$

$$X_2(dH)N_1 - C_1 - X_2^2(dH) = 0 \quad (9c)$$

$$X_2(dH)N_2 - C_2 - X_2 X_1(dH) = 0 \quad (9d)$$

Now, we put (8e), (8f) in (8b) to get:

$$\begin{aligned} & A_1 (X_1(dH)X_2 - X_2(dH)X_1) + A_2X_2 = \\ & -X_1(dH)(E_1(X_1(dH)X_2 - X_2(dH)X_1) + E_2X_1) - X_3X_1(dH)X_2 - \\ & X_2(dH)(D_1(X_1(dH)X_2 - X_2(dH)X_1) + D_2X_2) - X_3X_2(dH)X_1 \end{aligned}$$

grouping the terms in this equation we get the expression:

$$\begin{aligned} & (-(X_1(dH))^2E_1 - X_3X_1(dH) - X_2(dH)D_1X_1(dH) - \\ & X_2(dH)D_2 - A_1X_1(dH) - A_2)X_2 + \\ & (X_1(dH)E_1X_2(dH) - X_1(dH)E_2 + (X_2(dH))^2D_1 - \\ & X_3X_2(dH) + A_1X_2(dH))X_1 \end{aligned}$$

which is zero. Again, the linear independence of the vector fields give us two equations:

$$\begin{aligned} & A_2 + A_1X_1(dH) + D_2X_2(dH) + D_1X_1(dH)X_2(dH) + E_1(X_1(dH))^2 + \\ & X_3X_1(dH) = 0 \end{aligned} \tag{10a}$$

$$\begin{aligned} & A_1X_2(dH) + D_1(X_2(dH))^2 - E_2X_1(dH) + E_1X_1(dH)X_2(dH) - \\ & -X_3X_2(dH) = 0 \end{aligned} \tag{10b}$$

The equations (9a-d) and (10a-b) are what we need. From (9a) and (9c) we get C_1, C_2 . Using these two functions we get, with the help of (9b) and (9d) the functions B_1, N_2 given in the lemma. From (10b) we get the function A_1 . We put this function in (10a) to get A_2 . Hence the lemma.●

The algebra given in the lemma 4 is such that its representations allow us to construct extensions for Poisson tensors. However, it is complicated, but fortunately, we can choose the functions N_1, E_1, E_2, D_1, D_2 , in such a way that the algebra becomes easier to treat.

THEOREM2.-Three linearly independent vector fields X_1, X_2, X_3 , and a fourth vector X_H constructed as the contraction with an exact 1-co-tensor of the 2-tensor $X_1 \wedge X_2$, allow us to construct two compatible Poisson tensors if they are the base of the algebra with commutation rules:

$$\begin{aligned}
[X_1, X_2] &= 0 \\
[X_3, X_1] &= X_1 - X_2 \\
[X_3, X_2] &= 0
\end{aligned}$$

besides, the hamiltonian H of the Poisson structure $X_1 \wedge X_2$ satisfies the second order, factorizable, differential equation: $X_1 X_2(dH) = 0$.

PROOF.- The idea here is, again, quite simple: the lemma4 give us a 3-dimensional algebra obtained with two hypothesis: one vector is of a form such that only three vectors are required, and we choose the coefficients in such a way that three commutation rules are identities. Hence, now we shall choose the remaining coefficients and we shall use a new condition.

The choice for the coefficients is:

$$\begin{aligned}
N_1 &= E_1 = E_2 = 0, \\
D_1 &= -\frac{1}{X_2(dH)}, \quad D_2 = -1 + \frac{X_1(dH)}{X_2(dH)}
\end{aligned}$$

and the new condition is:

$$N_2 = 0 = (X_1 X_2 + X_2 X_1)(dH)$$

From this condition, and because $[X_1, X_2] = 0$, for the Hamiltonian we get the equation : $2X_1 X_2(dH) = 0$. Choosing the coefficients in the way which we have indicated reduce the algebra given in the lemma4 to the algebra given in the theorem 2. Hence, the theorem 2 is proved. •

We must remark that the other functions $A_1, A_2, B_1, B_2, C_1, C_2$ do not give us any condition, because we have not imposed any on them. Well, in fact there are many ways in which we can choose the functions, as must be clear, the one which we offer here is used because it is very easy to construct the representations in terms of first order differential operators, as we shall see in the example. As many possibilities are open, the notation for the algebra of the lemma 4 could be: $\mathfrak{S}(N_1, E_1, E_2, D_1, D_2)$, so, the case which we shall treat is: $\mathfrak{S}\left(0, 0, 0, -\frac{1}{X_2(dH)}, -1 + \frac{X_1(dH)}{X_2(dH)}\right) =_{def} \Delta$. The treatment which we have given is useful for Poisson tensors of the monomial form $X_1 \wedge X_2$, but our aim is not a general method to extend arbitrary Poisson tensors, instead, we are trying to construct singular-quasi-bi-Hamiltonian systems,

as we shall make in the next section. It is very important to remark that our algebraic criterion is sufficient, for this reason the consideration of the necessary conditions is not so important.

4 The quasi-bi-hamiltonian system

With our decomposable Poisson tensor, $\wp = X_1 \wedge X_2 + X_H \wedge X_3$, available through an algebraic criterion, is the moment of constructing the quasi-bi-Hamiltonian system. For this end we will use the infinitesimal automorphisms of the Poisson structures, which are given by:

$$X_H = dH \rfloor (X_1 \wedge X_2) = X_1 (dH) X_2 - X_2 (dH) X_1, \quad (11a)$$

$$X_F = dF \rfloor (X_H \wedge X_3) = X_H (dF) X_3 - X_3 (dF) X_H. \quad (11b)$$

Hence a relation between the different Hamiltonian vector fields is:

$$X_F = \{H, F\} X_3 + \rho(F) X_H, \quad \rho(F) = -X_3(dF). \quad (12)$$

So, if we choose F as an integral, or even as a Casimir, of the first Poisson tensor (Brouzet et al. only used the integrals, because the Casimir functions are not allowed for them), we easily get:

$$X_F = \rho X_H = \rho (dH \rfloor J), \quad (13a)$$

$$X_F(dF) = 0 \quad (13b)$$

$$X_F(dx^i) = \rho X_H(dx^i) \quad (13c)$$

As required by the definition (3). This procedure is, of course, suggested by Brouzet et al.[2][p. 2070-2071 eq. (4)]. Hence we have constructed an exact quasi-bi-Hamiltonian system on the basis of a decomposable Poisson tensor. For our quasi-bi-Hamiltonian dynamical system the property of being Pfaffian is not available, because this is defined by Brouzet et al. in connection with the Nijenhuis operator[2]. Now, we must remark here that the relations (11a-b) and the reduction to the form (13a) are the main motivations for the use of the suppositions (A) and (B) of the former section.

In this way we have, in general, constructed a quasi-bi-hamiltonian system on the basis of the algebra $\mathfrak{S}(N_1, E_1, E_2, D_1, D_2)$. We can see that, for example, the determination of the first Hamiltonian in terms of the available

elements, the vector fields, is possible in the case of the algebra Δ , and of course, if we have the form of this Hamiltonian, the second Hamiltonian is in principle, known.

Hence we have the family of QBH :

$$QBH_p = \langle M, X_1 \wedge X_2, H, X_H \wedge X_3, F \rangle \quad (14)$$

where in the last coordinate, instead of the function ρ , we use the second Hamiltonian because it is now available. There is a possible confusion here as F is, in fact, local, due to its definition as an integral of the first Poisson tensor (we mean: it is the solution of a partial differential equation of first order, and we understand this solution in the local sense). However, it is possible to find it explicitly in closed form in some simple cases. We will, nevertheless, restrict ourselves to this case. The algebra Δ is clearly solvable.

We can note an interesting fact of the algebra Δ : if $X_2 = 0$, then the differential equation: $\dot{x}_i = X_1(dx_i)$ by Lie theorem is integrable by quadratures[16], if it is 2-dimensional. Now, let us establish the connection with the Hojman method to construct Hamiltonian theories for autonomous first-order differential systems[1].

For the n -dimensional case consider the equation: $\dot{x}_i = X_1(dx_i)$. This differential equation admits a Poisson structure of the form $J = X_1 \wedge X_3$, by just choosing the Hamiltonian as $X_1(dH) = 0$ (clearly a particular case of the differential equation: $X_1X_2(dH) = X_2X_1(dH) = 0$ and $X_1(dH) = cte.$) because by contraction we get the vector field: $X_3(dH)X_1$. Under this condition $\rho = X_3(dH)$ is a constant of the motion (PROOF: $X_3(X_1(dH)) - X_1(X_3(dH)) = X_1(dH) \Rightarrow X_1(X_3(dH)) = 0$) as required by Hojman method. Hence the scaling used by Hojman (see ref.[1] p. 669 equation (12)) for the vector field $\bar{X}_1 = \rho X_1$ (in components of the vector fields it is: $\bar{\eta}^i = \rho \eta^i$) is possible. For this reason the Hamiltonian structure is for the system with vector field of the form: $\bar{X}_1 = X_1 \rho$ which can be seen as Hamiltonian. The Hamiltonian vector field constructed by Hojman is, thus: $dH \rfloor J = \bar{X}_H = \rho X_1$. For this reason, and because $X_2 = 0$ can be seen as a particular case of the algebra Δ . Then, our construction covers that one of Hojman.

This is the unique way in which the connection can be established, because it is possible to commit the mistake of trying to apply the procedure which we offer to construct an extended Poisson tensor over that constructed by the method of Hojman. If we do this we get a contradiction. Let us show this:

Let $X_1 \wedge X_3$ be the Poisson tensor obtained by the Hojman method, then, our procedure gives us the extension with the help of the tensor: $\bar{X}_H \wedge Y$, and the conditions:

$$\begin{aligned} [Y, X_1] &= X_1 - X_3, \\ [X_1, X_3] &= 0, \\ [X_3, Y] &= 0. \end{aligned}$$

But, according to the Hojman method we must have: $[X_3, X_1] = X_1$, which cannot be satisfied if we use the methodology proposed in the method which we offer. Hence, this procedure is wrong.

So, we reject any criticism to our method on the basis of the preceding calculation, which comes from an incorrect consideration. The only way to establish the connection is that which considers the Hojman method as a particular case of ours in the lines which we have already explained. Let us make a last comment: Hojman proposed in his paper[1] the idea that if the differential equation in question has, say, m -symmetries ($m \leq n$), then, on the basis of his method, entirely based on the assumption $X_2 = 0$ for the algebra Δ , we can get new Poisson tensors. So, we must find a representation for the algebra:

$$[X_i, X_1] = X_1, i = 1, \dots, m. \quad (15)$$

Hence, if $m = n$ we get the result (not remarked in Hojman paper) that any n -dimensional integrable system: $\dot{x}_i = X_1(dx_i)$, admits n Poisson structures. The vector fields X_i are of course, the classical Lie symmetries of the differential equation. In fact, Hojman proposed (ref [1] p. 673) a way to extend the Poisson tensor constructed by his method, but based on the relation (15) and, because this relation is deduced from our algebra Δ , the Poisson tensor extended by Hojman method are compatible.

If we want to get a singular bi-Hamiltonian system for the 2-dimensional case (1-dimensional if we use Darboux coordinates) we must add the condition: $X_3(dF) = -1$, to the condition $\{H, F\} = 0$, to get the second hamiltonian F , which shows that this case is more restricted, from the point of view of our calculations, than the singular one.

It is important to remark two points:

(1).- The method which we propose to construct decomposable Poisson tensors is, essentially, a problem of algebra representations[3], because any

realization of the Lie algebra Δ can be used to construct a decomposable Poisson tensor. The infinitesimal automorphisms which leave invariant our decomposable Poisson tensor $\wp$ are generated, as before, by the vector fields: $X_\gamma = \gamma \rfloor \wp$ with γ any 1-form.

(2).-If we choose such vectors so that the commutation relations which define $\mathfrak{S}$ are not realized, our Poisson tensors are not compatible and thus, we do not have the possibility of reducing the full algebra to Δ which gives us the operative formulation for the construction of a quasi-bi-Hamiltonian dynamical system.

5 Digression on Jacobi structures.

It is not our aim to discuss in full detail the Jacobi structures[9][8], but we think that it is important to remark how the methodology which we propose for the extension of Poisson tensors could be useful for constructing Jacobi structures.

We start with a general definition (see ref. [14] p. 6314):

DEFINITION 5.- On a smooth (C^∞) manifold M we say that we have a Jacobi structure if, and only if, it is possible to construct a pair $\langle \Lambda, X_H \rangle$ such that Λ is a 2-contraction and X_H is 1-contraction on M such that the following conditions holds:

$$[\Lambda, \Lambda] = 2X_H \wedge \Lambda, [X_H, \Lambda] = 0 \quad (16)$$

We use the notation X_H to connect with the former sections. Clearly, this is not a Hamiltonian vector field unless otherwise stated, it is an arbitrary vector field. Our methodology is summarized in the following:

THEOREM 3.- Three linearly independent vector fields X_1, X_2, X_H , allows us to construct a Jacobi structure of the form $\langle X_1 \wedge X_2, X_H \rangle$ on a smooth manifold M if they satisfy the commutation rules:

$$[X_1, X_2] = -X_H \quad (16a)$$

$$[X_H, X_1] = -AX_1 + BX_2 \quad (16b)$$

$$[X_H, X_2] = CX_1 + AX_2 \quad (16c)$$

with A, B, C , arbitrary functions.

PROOF: The assertion in the theorem just requires the sufficiency of the commutation rules (16a-c) to be shown, that is: if the vector fields form the algebra with commutation rules (16a-c), which we shall denote as $\mathcal{J}(A, B, C)$, then we can construct the pair $\langle X_1 \wedge X_2, X_H \rangle$. The proof is as follows: expand the Schouten bracket to get the equation:

$$2X_1 \wedge [X_1, X_2] \wedge X_2 = 2X_H \wedge (X_1 \wedge X_2)$$

so, according to the definition we must have:

$$[X_1, X_2] = -X_H$$

the other condition simply states that the vector field X_H is an infinitesimal automorphism of the 2-contr-tensor. We meet sufficient conditions for this requirement in lemma2, the conditions are equal to (16a-b), but we changed the arbitrary functions to avoid confusion. Hence, if these algebraic conditions are satisfied, we can construct a Jacobi structure on the manifold, by the conditions in the definition. The theorem is proved.●

We shall give 1 example below.

6 Examples.

The key points for constructing an example of an exact quasi-bi-Hamiltonian system are: a Poisson tensor, its contraction with an integrable 1-form and an arbitrary vector. The main source of all the examples is the work of Cariñena et al.[3] on solvable Lie algebras.

We will consider three ortogonal vectors, for the sake of generality and non-triviality. In this case the algebra Δ to be satisfied is:

$$\begin{aligned} [X_1, X_2] &= 0 \\ [X_3, X_1] &= X_1 - X_2 \\ [X_3, X_2] &= 0 \end{aligned}$$

In each example we will use a super-index to denote the particular realization of the algebra and the number of the example.

1.- Every 2-dimensional example is trivial, because the Schouten brackets are 3-tensors which vanish in two dimensions, by construction.

2.-Consider a 3-dimensional (the first non-trivial dimension for a 3-vector like the Schouten bracket) example with the Poisson tensor:

$$J^{(2)} = (x_1 \frac{\partial}{\partial x_2} - x_2 \frac{\partial}{\partial x_1}) \wedge \frac{\partial}{\partial x_3}, \quad (21a)$$

$$X_1^{(2)} = x_1 \frac{\partial}{\partial x_2} - x_2 \frac{\partial}{\partial x_1}, X_2^{(2)} = \frac{\partial}{\partial x_3} \quad (21b)$$

and the arbitrary vector:

$$X_3^{(2)} = P_1(x_1, x_2) \frac{\partial}{\partial x_1} + P_2(x_1, x_2) \frac{\partial}{\partial x_2} + P_3(x_1, x_2) \frac{\partial}{\partial x_3}. \quad (22)$$

Clearly $X_1^{(2)}$, $X_2^{(2)}$, $X_3^{(2)}$ satisfy the required commutation relations. The other commutation relation gives us the partial differential equations:

$$\begin{aligned} x_2 \frac{\partial P_1}{\partial x_1} - x_1 \frac{\partial P_1}{\partial x_2} - P_2 &= x_2, \\ x_2 \frac{\partial P_2}{\partial x_1} - x_1 \frac{\partial P_2}{\partial x_2} + P_1 &= -x_1, \\ x_2 \frac{\partial P_3}{\partial x_1} - x_1 \frac{\partial P_3}{\partial x_2} &= 1, \end{aligned}$$

for the unknown functions P_1, P_2, P_3 . The solution for these equations defines the required vector to get the new Poisson structure to construct the decomposable tensor. We can re-write the equations as:

$$(x_2 \frac{\partial}{\partial x_1} - x_1 \frac{\partial}{\partial x_2})^2 P_1 + P_1 = 0, \quad (23a)$$

$$(x_2 \frac{\partial}{\partial x_1} - x_1 \frac{\partial}{\partial x_2}) P_1 - x_2 = P_2 \quad (23b)$$

$$(x_2 \frac{\partial}{\partial x_1} - x_1 \frac{\partial}{\partial x_2}) P_3 = 1. \quad (23c)$$

So, we just need to solve the equations (21a, 21c). We can solve this system with the help of the transformation to polar coordinates: $x_1 = r \cos \theta$, $x_2 = r \sin \theta$, to get the equations:

$$\begin{aligned}
\frac{\partial^2}{\partial \theta^2} p_1(r, \theta) + p_1(r, \theta) &= 0, \\
\frac{\partial}{\partial \theta} p_1(r, \theta) - r \sin \theta &= p_2(r, \theta) \\
\frac{\partial}{\partial \theta} p_3(r, \theta) &= 1.
\end{aligned}$$

With $P_j(x_1(r, \theta), x_2(r, \theta)) = p_j(r, \theta)$. Hence, a solution is:

$$\begin{aligned}
P_1 &= A(x_1^2 + x_2^2) \sin(\arctan(\frac{x_2}{x_1}) + B(x_1^2 + x_2^2)), \\
P_2 &= A(x_1^2 + x_2^2) \cos(\arctan(\frac{x_2}{x_1}) + B(x_1^2 + x_2^2)) - x_2, \\
P_3 &= \arctan(\frac{x_2}{x_1}) + C(x_1^2 + x_2^2).
\end{aligned}$$

This is enough to construct the second Poisson tensor which makes the extension of $J^{(2)}$ and the particular $QBH_{(2)}^*$, like in the first example, by just choosing a generator H as solution of its corresponding partial differential equation. The calculation of the extended Poisson tensor gives us:

$$\begin{aligned}
\wp^{(1)} &= (x_1 \frac{\partial}{\partial x_2} - x_2 \frac{\partial}{\partial x_1}) \wedge \frac{\partial}{\partial x_3} + \\
& ([x_1 \frac{\partial H^{(2)}}{\partial x_2} - x_2 \frac{\partial H^{(2)}}{\partial x_1}] \frac{\partial}{\partial x_3} - \frac{\partial H^{(2)}}{\partial x_3} [x_1 \frac{\partial}{\partial x_2} - x_2 \frac{\partial}{\partial x_1}]) \wedge \\
& (A(x_1^2 + x_2^2) [\sin(\arctan(\frac{x_2}{x_1})) \frac{\partial}{\partial x_1} + \cos(\arctan(\frac{x_2}{x_1})) \frac{\partial}{\partial x_2}] + \\
& + \arctan(\frac{x_2}{x_1}) \frac{\partial}{\partial x_3} + C(x_1^2 + x_2^2) \frac{\partial}{\partial x_3}).
\end{aligned}$$

In this case the Hamiltonian of the first Poisson structure must be a solution of the differential equation:

$$x_1 \frac{\partial^2 H^{(2)}}{\partial x_2 \partial x_3} - x_2 \frac{\partial^2 H^{(2)}}{\partial x_1 \partial x_3} = 0$$

3.- Consider now the case of linear Poisson tensors: $J = X_A \wedge X_a$, which define semi-direct extensions of Abelian Lie algebras[3] (the two preceding examples are particular cases of this):

$$\begin{aligned}
X_1 &= X_A = \sum_{i,j=1}^{n-1} A_i^j x_j \frac{\partial}{\partial x_i}, \\
X_2 &= X_a = \frac{\partial}{\partial x_n}, \\
X_3 &= \sum_{j=1}^n P_j(x_1, \dots, x_{n-1}) \frac{\partial}{\partial x_j},
\end{aligned}$$

when they act on functions belonging to $C^\infty(M, \mathfrak{R})$; which is the case important for us now. It is clear that we will take the tensor $J = X_A \wedge X_a$ as our initial tensor. We can see that our vector fields satisfy the required commutation relations if the following set of first order partial differential equations (obtained after a straightforward calculation which we can omit here) for the functions $P_j(x_1, \dots, x_{n-1})$ is solvable:

$$\begin{aligned}
\sum_{k=1}^{n-1} A_k^j x_j \frac{\partial P_i}{\partial x_k} &= A_i^j (P_j - x_j), \forall (i, j) : i, j \in \{1, \dots, n-1\}, \\
\left(\sum_{i,j}^{n-1} A_i^j x_j \frac{\partial}{\partial x_i} \right) P_n &= 1.
\end{aligned}$$

From a theoretical point of view, an analytic solution for this set of differential equations exists by the well-known Cauchy-Kovalevskaya theorem[17]. Hence, there is always (at least locally) an extension to the Poisson tensor which defines semi-direct extensions of abelian Lie algebras on the basis of the method which we propose (of course, for the use of the Cauchy-Kovalevskaya theorem we must change $C^\infty(M, \mathfrak{R})$ by $C^\omega(M, \mathfrak{R})$, the Banach space of analytic functions). A classification of these algebras in dimension 1, 2, 3, 4 is given by Cariñena et al[3]. and for all the Poisson tensors which they consider is valid the methodology which we propose, with just one remark: the Poisson tensor must be of the form $X_1 \wedge X_2$. Our method of extension is not the same as the one of these authors, although we use a common property: in the second Poisson tensor $X_H \wedge X_3$, we take X_H as a derivation of the first Poisson tensor. But this is all, because they want to get all the derivation algebra, whereas we fix a derivation; a Hamiltonian vector field as derivation; and construct the remaining piece: the vector field X_3 . So, if we know the

algebra of derivations, our method can be applied to each element of this algebra to get a different extension for each one of its elements.

4.- Let us show that $j(0, -1, -1) \simeq so(3)$. The commutation rules for $so(3)$ are well-known to be (we use the circumflex accent to avoid any confusion with the manifold coordinates):

$$[\hat{x}_1, \hat{x}_2] = \hat{x}_3, [\hat{x}_2, \hat{x}_3] = \hat{x}_1, [\hat{x}_3, \hat{x}_1] = \hat{x}_2$$

while those rules for the algebra $j(0, -1, -1)$ are:

$$[X_1, X_2] = -X_H, [X_H, X_1] = -X_2, [X_2, X_H] = -X_1$$

the isomorphism is, clearly, the linear map: $X_H = -\hat{x}_3$, $X_2 = \hat{x}_2$, $X_1 = \hat{x}_1$. So, any representation of $so(3)$ can be used to get Jacobi structures, for example, the usual one of the form:

$$X_H = x_2 \frac{\partial}{\partial x_1} - x_1 \frac{\partial}{\partial x_2}, \quad X_2 = x_3 \frac{\partial}{\partial x_1} - x_1 \frac{\partial}{\partial x_3}, \quad X_3 = x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2}$$

the Jacobi structure for this case is $\langle X_1 \wedge X_2, X_H \rangle$

COMMENT.-Well, in fact every 3-dimensional Lie algebra can be classified by means of linear transformations in the manifold coordinates:

$$X_i = \sum_j a_{ij} \hat{X}_j$$

where we suppose that the coefficients are just constants (hence we left out non-linear transformations of the coordinates by general diffeomorphisms). The conditions which we must impose on the linear transformation are: (1).- it is an isomorphism, we mean, a bijective homomorphism of Lie algebras, (2).- the Jacobi identity. This has been done, for example, by Bryant[7] [p.37-40] and we refer there for further details on the reduction.

7 Conclusions.

We have shown how to construct extensions of Poisson tensors and singular-quasi-bi-Hamiltonian systems on this basis. The procedure seems much more comprehensive than the method due to Brouzet et al. although it is valid only for those Poisson tensors of the form: $X_1 \wedge X_2$ and its contractions over sets of integrable (exact) 1-forms. In principle it is possible

to construct as many quasi-bi-hamiltonian systems as smooth scalar generators are imaginable, with just one constraint; let us show why. A solution for the partial differential equation $X_1X_2(dH) = 0$ can be taken as $H = I_1(\xi_1) + I_2(\xi_2)$ where each ξ_i is an invariant function of each vector field X_i . Hence, by the commutativity of the fields this is a solution: $X_1X_2(I_1) + X_1X_2(I_2) = X_2X_1(I_1) + X_1X_2(I_2) = 0$, because $X_1(I_1) = 0$, $X_2(I_2) = 0$, and because the functions I_1, I_2 are arbitrary, our assertion is justified. The constraint is, clearly, to move only along the characteristic paths defined by the invariants.

We can see one case in which the Hamiltonian will be totally arbitrary: if the vector fields X_1, X_2 commute and anti-commute the first Hamiltonian is arbitrary, there is no restriction on its form. However we can see that the conditions: $X_1X_2 = X_2X_1$ $X_1X_2 = -X_2X_1$ are fulfilled if we suppose that $X_1X_2 = 0$ an operator identity. But this is our condition to get the first Hamiltonian, hence nothing new arises.

As a second point, we see that the method of extension is different from that due to Cariñena et al.[3], because, as explained in the example 3, the method can be applied to each element of the derivation algebra to get different extensions of the same Poisson tensor.

Acknowledgments.

This work has been supported in part by the financial aid of CONACYT Research Project 33471-E and the Centro de Estudios Multidisciplinarios UAZ. RAF and MAG are grateful to Ana Maria Damore for her patient to read the manuscript and to Rogelio Cárdenas Hernández by his interest in the development of the work. MAG acknowledges to Professor V. G. Makhankov for his constant support discussions.

References

- [1] Hojman, S. A.: The construction of a Poisson structure out of a symmetry and a conservation law.
J. Phys. A: Math. Gen. v.29 n.3 (1996) p. 667-674.
- [2] Brouzet, R., R. Caboz, J. Rabenivo and V. Ravoson: Two degrees of freedom quasi-bi-Hamiltonian systems.
J. Phys. A: Math. Gen. v.29 n. 9 (1996) p. 2069-2076.

- [3] Cariñena, J.F., A. Ibort, G. Marmo and A. Perelomov: On the geometry of Lie algebras and Poisson tensors.
J. Phys. A: Math Gen. v. 27 n. 22 (1994) p. 7425-7449.
- [4] Bueken, P.: Multi-Hamiltonian structures for a class of degenerate completely integrable systems.
J. Math. Phys. v. 37 n. 6 (1996) p. 2851-2862.
- [5] Goedert J., F. Haas, D. Hua, M. R. Feix and L. Cairó: Generalized constant structures for systems in three dimensions with a rescalable constant of motion.
J. Phys. A: Math. and Gen. v.27 n19 (1994) p.6495-6507
- [6] Jost, Res : Poisson Brackets (An unpedagogical Lecture).
Rev. Mod. Phys. April (1964). p. 572-579.
- [7] Bryant, R.L.: An introduction to Lie Groups and Symplectic Geometry. (Lecture 2 passim 37-40)
Geometry and Quantum Field Theory, D.S. Freed, K.K. Uhlenbeck (eds.). AMS (1995)
- [8] Lichnerowicz, A.: Les variétés de Poisson et les algèbres de Lie associées.
J. Diff. Geom. v. 12 (1977), p. 253-300.
- [9] Kirillov, A.: Local Lie Algebras.
Russian Math Surveys v. 31 (1976) p. 55-75.
- [10] Weinstein, A.: The local structure of Poisson manifolds.
J. Diff. Geom. v. 18 (1988 p. 523-557).
- [11] Kostant, B.: Orbits, symplectic structures and representation theory.
Proc. US-Japan. Seminar on Diff. Geom., Kyoto. Nippon Hyronsha, Tokyo 77 (1966).
- [12] Souriau, J. M.: Structure des Systèmes Dynamiques.
Dunod, Paris. (1970).

- [13] Gillemin, V. and S. Sternberg: Symplectic Techniques in Physics.
Cambridge University Press (1984).
- [14] Chinea, Domingo, Juan C. Marrero and Manuel de León: Prequantizable Poisson manifolds and Jacobi structures.
J. Phys. A: Math and Gen v 29 (1996) p 6313-6324.
- [15] Marsden, J.E. and T. S. Ratiu: Introduction to Mechanics and Symmetry.
Texts in Applied Mathematics n.17 (1994) Springer-Verlag, New York.
- [16] Arnold, V.I. (ed.): Dynamical Systems III.
Encyclopaedia of Mathematical Sciences v. 3 (1993) Springer-Verlag, Berlin.
- [17] Courant, R. and D. Hilbert : Methods of Mathematical Physics. Volume II.
Wiley Classics Edition (1989). Interscience Publishers, New York.